

*Journal of Agricultural Science* (2013), **151**, 154–162. doi:10.1017/S0021859612000263

© Cambridge University Press 2012. The online version of this article is published within an Open Access environment subject to the conditions of the Creative Commons Attribution-NonCommercial-ShareAlike licence <<http://creativecommons.org/licenses/by-nc-sa/2.5/>>. The written permission of Cambridge University Press must be obtained for commercial re-use.

## CLIMATE CHANGE AND AGRICULTURE RESEARCH PAPER

# Effects of elevated CO<sub>2</sub> and temperature on seed quality

J. G. HAMPTON<sup>1</sup>\*, B. BOELT<sup>2</sup>, M. P. ROLSTON<sup>3</sup> AND T. G. CHASTAIN<sup>4</sup>

<sup>1</sup> *Bio-Protection Research Centre, Lincoln University, Lincoln 7647, New Zealand*

<sup>2</sup> *Sciences and Technology, Aarhus University, DK 4200 Slagelse, Denmark*

<sup>3</sup> *AgResearch Ltd, Lincoln 7647, New Zealand*

<sup>4</sup> *Department of Crop and Soil Science, Oregon State University, Corvallis, OR 97331-3002, USA*

(Received 30 August 2011; revised 23 November 2011; accepted 29 February 2012;  
first published online 30 March 2012)

### SUMMARY

Successful crop production depends initially on the availability of high-quality seed. By 2050 global climate change will have influenced crop yields, but will these changes affect seed quality? The present review examines the effects of elevated carbon dioxide (CO<sub>2</sub>) and temperature during seed production on three seed quality components: seed mass, germination and seed vigour.

In response to elevated CO<sub>2</sub>, seed mass has been reported to both increase and decrease in C<sub>3</sub> plants, but not change in C<sub>4</sub> plants. Increases are greater in legumes than non-legumes, and there is considerable variation among species. Seed mass increases may result in a decrease of seed nitrogen (N) concentration in non-legumes. Increasing temperature may decrease seed mass because of an accelerated growth rate and reduced seed filling duration, but lower seed mass does not necessarily reduce seed germination or vigour.

Like seed mass, reported seed germination responses to elevated CO<sub>2</sub> have been variable. The reported changes in seed C/N ratio can decrease seed protein content which may eventually lead to reduced viability. Conversely, increased ethylene production may stimulate germination in some species. High-temperature stress before developing seeds reach physiological maturity (PM) can reduce germination by inhibiting the ability of the plant to supply the assimilates necessary to synthesize the storage compounds required for germination.

Nothing is known concerning the effects of elevated CO<sub>2</sub> on seed vigour. However, seed vigour can be reduced by high-temperature stress both before and after PM. High temperatures induce or increase the physiological deterioration of seeds. Limited evidence suggests that only short periods of high-temperature stress at critical seed development stages are required to reduce seed vigour, but further research is required.

The predicted environmental changes will lead to losses of seed quality, particularly for seed vigour and possibly germination. The seed industry will need to consider management changes to minimize the risk of this occurring.

### INTRODUCTION

With global change, atmospheric carbon dioxide (CO<sub>2</sub>) concentration is predicted to rise from today's value of c. 370–550 ppm by 2050 and could reach between 730 and 1010 ppm by 2100 (Solomon *et al.* 2007). This, combined with other atmospheric changes, is projected to increase global mean temperatures by 1.4–5.8 °C (Houghton *et al.* 2001). Jaggard *et al.* (2010) concluded that CO<sub>2</sub> enrichment was likely to allow yield increases of c. 13% in most C<sub>3</sub>

crops, but yields of C<sub>4</sub> crops are not expected to change. However, increasing temperatures may negate these increases in C<sub>3</sub> crops, particularly if they occur during reproductive growth (Allen & Boote 2000; Wheeler *et al.* 2000). Gornall *et al.* (2010) noted that extreme weather events are more likely to occur in the changed climate of the future, and predicted that over much of the world's crop land, maximum daily temperature highs may be increased by around 3 °C by 2050.

A major challenge ahead for those involved in the seed industry, therefore, is to provide cultivars that can maximize future crop production in a changing

\* To whom all correspondence should be addressed. Email: john.hampton@lincoln.ac.nz

climate (Ainsworth *et al.* 2008a; Bruins 2009; Ceccarelli *et al.* 2010). Ainsworth *et al.* (2008b) considered that this will be possible within a decade.

Successful crop production in any environment depends initially on the quality of the seed being sown. The term 'seed quality' is used in practice to describe the overall value of a seed lot for its intended purpose (Hampton 2002), and includes the components of species and cultivar purity, seed mass (size), physical purity, germination, vigour, moisture content and seed health. The present review examines the effects of increased CO<sub>2</sub> and increased temperature on three of these seed quality components, seed mass, germination and vigour.

## SEED MASS

Within the seed industry, seed size is commonly denominated by the mean seed weight, often expressed as 'thousand seed weight', the weight of 1000 seeds of the seed lot. However, seed size refers to volume, while seed weight and seed mass refer to density, which are different traits (Castro *et al.* 2006). Seed mass in crop cultivars is considered the least variable of the seed yield components because of plant breeding for increased seed uniformity (Almekinders & Louwaars 1999) and the removal of small seeds during the seed cleaning processes. Factors affecting seed mass, including genetic factors, water availability and nutrient availability were reviewed by Castro *et al.* (2006).

Increased atmospheric CO<sub>2</sub> concentrations might be expected to increase seed mass because of increased plant assimilate availability (Jablonski *et al.* 2002), but the reported effects of elevated CO<sub>2</sub> are highly variable among species (Miyagi *et al.* 2007; Hikosaka *et al.* 2011). Different studies have reported seed mass to increase (Musgrave *et al.* 1986; Baker *et al.* 1989; Dijkstra *et al.* 1999; Steinger *et al.* 2000; Quaderi & Reid 2005), show no change (Edwards *et al.* 2001; Prasad *et al.* 2002) and decrease (Huxman *et al.* 1998; Smith *et al.* 2000; Wagner *et al.* 2001) in response to elevated CO<sub>2</sub>. Jablonski *et al.* (2002) conducted a meta-analysis of 184 CO<sub>2</sub> enrichment studies from 79 species and found a mean 4% increase in seed mass, with the response being greater in legumes (+8%) than non-legumes (+3%), and absent in C<sub>4</sub> plants. Considerable variation in seed mass in response to elevated CO<sub>2</sub> was also reported within species. Hikosaka *et al.* (2011) reported the enhancement ratio of seed mass per plant (seed mass in

elevated CO<sub>2</sub>/seed mass in ambient CO<sub>2</sub>) ranged from 0.75 to 4.45 in rice (*Oryza sativa* L.), from 0.93 to 1.87 in soybean (*Glycine max* (L.) Merrill), and from 0.88 to 2.07 in wheat (*Triticum aestivum* L.).

Jablonski *et al.* (2002) found a 14% reduction in seed nitrogen (N) in response to elevated CO<sub>2</sub> averaged across all 79 species in their analysis, although there was significant variation; seed N was not reduced in legumes, but was reduced in non-legumes. Hikosaka *et al.* (2011) suggested that seed mass could only increase when N became more available at elevated CO<sub>2</sub> concentrations. Legumes may use increased carbon (C) gain under elevated CO<sub>2</sub> for increased nitrogen fixation (Allen & Boote 2000), and can therefore increase seed mass without decreasing seed N. In non-legumes, seed mass increases may result in a decrease in seed N concentration. In some species, this decreased seed N may be at the expense of seed quality (Fenner 1991; Andalo *et al.* 1996).

Increasing temperature can negate the response to elevated CO<sub>2</sub> (Prasad *et al.* 2002) and may reduce seed mass (Spears *et al.* 1997) because of the resulting acceleration in seed growth rate (dry matter accumulation) and reduction in the duration of seed filling (Weigand & Cueller 1981; Young *et al.* 2004). However, a reduction in the rate of seed dry matter accumulation can also occur (Gibson & Paulsen 1999) and seed mass has also been reported not to change, or sometimes increase, in response to temperature increase (Peltonen-Sainio *et al.* 2011). A reduced seed mass for a seed lot does not necessarily mean a loss in other seed quality attributes. Many studies have shown no relationship between seed mass and germination (Castro *et al.* 2006) or seed mass and seed vigour (Powell 1988).

## GERMINATION

For high-quality seed lots, germination (defined as the process that begins with imbibition and which is completed by the production of a normal seedling; ISTA 2011) is desired by the seed industry to be as close to 100% as possible. The germination of a seed lot can be negatively affected by the conditions the seeds are exposed to during harvesting, drying, cleaning and storage, but can also be reduced by unfavourable environmental conditions in the field during seed growth and development (Dornbos 1995), particularly temperature, rainfall and relative humidity (Egli *et al.* 2005).

Table 1. *Effect of temperature during seed development on seed germination and seed vigour of two soybean cultivars (adapted from Spears et al. 1997)*

|  | Temperature regime and duration* |       |          |       |          |       |
|--|----------------------------------|-------|----------|-------|----------|-------|
|  | 27/22 °C                         |       | 33/28 °C |       | 38/33 °C |       |
|  | R5-PM                            | R5-R8 | R5-PM    | R5-R8 | R5-PM    | R5-R8 |
| Germination (%)                            |                                  |       |          |       |          |       |
| McCall†                                    | 100                              | 100   | 98       | 100   | 94       | 63    |
| Hutchenson                                 | 100                              | 100   | 99       | 98    | 57       | 14    |
| Germination (%) after accelerated ageing*‡ |                                  |       |          |       |          |       |
| McCall†                                    | 98                               | 100   | 73       | 86    | 8        | 7     |
| Hutchenson                                 | 98                               | 95    | 43       | 86    | 2        | 1     |
| Conductivity‡ (µS/cm/g)                    |                                  |       |          |       |          |       |
| McCall†                                    | 5.5                              | 5.4   | 7.0      | 6.3   | 16.1     | 23.9  |
| Hutchenson                                 | 6.5                              | 6.2   | 9.1      | 10.6  | 29.8     | 35.9  |

\* Day/night temperatures with 10 h at the day temperature; R5=beginning of seed fill; PM=physiological maturity; R8=harvest maturity.

† Soybean cultivars; McCall=indeterminate growth habit; Hutchenson=determinate growth habit.

‡ Seed vigour tests.

Seed germination in response to elevated CO<sub>2</sub> has been reported to decrease (Farnsworth & Bazazz 1995; Andalo *et al.* 1996; Quaderi & Reid 2005), show no change (Huxman *et al.* 1998; Steinger *et al.* 2000; Thomas *et al.* 2009; Way *et al.* 2010) or increase (Wulf & Alexander 1985; Ziska & Bunce 1993; Edwards *et al.* 2001). The responses vary among species (Ziska & Bunce 1993) and genotypic variation has also been reported (Andalo *et al.* 1996).

Elevated CO<sub>2</sub> has been shown to increase the C/N ratio in seeds (Huxman *et al.* 1998; Steinger *et al.* 2000; He *et al.* 2005) and in non-legumes, seed N reduction can occur when seed mass is increased by elevated CO<sub>2</sub> (see previous section). High seed N is an advantage for germination rate (Hara & Toriyama 1998), but not germination *per se*. However, a change in C/N ratio can lead to a decrease in seed protein content, resulting in a reduction in the ability of the seed to supply the amino acids required for the *de novo* protein synthesis necessary for embryo growth in the germinating seed. This could reduce seed viability (Andalo *et al.* 1996).

Elevated CO<sub>2</sub> also increases ethylene production (Esashi *et al.* 1986) and Ziska & Bunce (1993) suggested that an increased availability of ethylene may have been the reason for the stimulated germination they reported. Ethylene is implicated in the promotion of germination of non-dormant seeds of many species (Leubner-Metzger 2006).

In different plant species, sometimes even small differences in temperature during seed development and maturation can have an influence on germination (Guterman 2000). High temperatures during seed filling frequently disrupt normal seed development, which increases the proportion of seeds that are shrivelled, abnormal and are of lower quality (Spears *et al.* 1997). However, it has been shown that after removal of these seeds, the germination of the remaining seeds decreases as mean maximum temperature during seed filling increases (Khalil *et al.* 2001, 2010; Egli *et al.* 2005; Thomas *et al.* 2009; Table 1). High-temperature stress before the developing seeds achieve physiological or mass maturity (PM – the end of the seed filling phase) is likely to inhibit the ability of the plant to supply the seeds with the assimilates necessary to synthesize the storage compounds required during the germination process (Dornbos & McDonald 1986), and/or the seeds suffer physiological damage (see McDonald & Nelson 1986; Coolbear 1995; Powell 2006) to the extent that the ability to germinate is lost.

High-temperature stress after PM can also sometimes reduce germination (Green *et al.* 1965; Table 1), but more often reduces seed vigour (see next section).

The relationship between temperature during seed development and subsequent seed germination requires further investigation. For example in soybean, temperatures (32–38 °C) that reduced the germination

Table 2. Effect of high temperature (30/25 °C) for 4 days at different stages of seed development and maturation in two cultivars of pea (*Pisum sativum* L.) on seed quality components (adapted from Shinohara et al. 2006b)

| Stage at treatment* | Mean seed weight (g) |            | Germination (percentage) |           | Hollow heart (proportion) |            | Average conductivity† (µS/cm/g) |           |
|---------------------|----------------------|------------|--------------------------|-----------|---------------------------|------------|---------------------------------|-----------|
|                     | Alderman‡            | E. Onward‡ | Alderman                 | E. Onward | Alderman                  | E. Onward  | Alderman                        | E. Onward |
| Control             | 379                  | 350        | 95                       | 93        | 0.04                      | 0.00       | 220                             | 363       |
| S1                  | 309                  | 289        | 84                       | 97        | 0.01                      | 0.03       | 315                             | 368       |
| S2                  | 326                  | 262        | 86                       | 92        | 0.24                      | 0.04       | 367                             | 374       |
| S3                  | 391                  | 345        | 86                       | 96        | 0.08                      | 0.00       | 339                             | 467       |
| S4                  | 378                  | 362        | 88                       | 98        | 0.01                      | 0.00       | 371                             | 475       |
| S5                  | 376                  | 337        | 92                       | 96        | 0.04                      | 0.00       | 424                             | 420       |
| S.E.D. (D.F.)       | 11 (18)              | 9 (18)     | 6.7 (18)                 | 3.9 (18)  | 0.054 (18)                | 0.020 (18) | 31 (140)                        | 123 (140) |

\* S1=beginning of seed filling (810 mg/g SMC); S2=rapid seed filling (700 mg/g SMC); S3=PM (630 mg/g SMC); S4=beginning of desiccation (440 mg/g SMC); S5=harvest maturity (230 mg/g SMC); SMCs are mean of the two cultivars.

† Data are the average of 25 results in the single seed conductivity vigour test.

‡ Pea cultivars.

S.E.D. (between cultivars)=10 (mean seed weight), 5.4 (germination), 0.041 (hollow heart) and 71 (average conductivity).

of some cultivars in controlled environments did not vary during seed filling, in contrast to field temperatures which can vary substantially (Egli *et al.* 2005), and the plants were at these temperatures from anthesis until seed harvest. However, there may be critical periods during seed development when seeds are particularly sensitive to temperature (Egli *et al.* 2005; Shinohara *et al.* 2006a). This was investigated for pea (*Pisum sativum* L.) by Shinohara *et al.* (2006a), who showed that when plants were exposed to a day/night temperature of 30/20 °C for 4 days (=240 °C h above a base temperature (T<sub>b</sub>) of 25 °C) at the beginning of seed filling and then returned to the field until seed harvest, germination was significantly reduced in one of two cultivars (Table 2). Exposure to these conditions at later stages of seed development did not affect germination.

## SEED VIGOUR

While the term germination has long been used to describe the planting value of a seed lot (ISTA 2011), when conditions in the seed bed are less than optimal the germination test is a poor predictor of field emergence (Dornbos 1995), suggesting that a further physiological aspect to seed quality exists—seed vigour (Powell 2006). Seed vigour is defined by ISTA (2011) as ‘the sum of those properties that determine the activity and level of performance of seed lots of acceptable germination in a wide range of

environments’, or more simply, the ability of a high germination seed lot to emerge under seed-bed stress.

While there have been reports that elevated CO<sub>2</sub> increases or decreases seedling vigour (i.e. growth rate or biomass production) because of the effect on seed mass (Huxman *et al.* 1998; Steinger *et al.* 2000), there have been no reports on the effects of elevated CO<sub>2</sub> on seed vigour.

Seed vigour is reduced by high-temperature stress before PM (Spears *et al.* 1997; Egli *et al.* 2005; Shinohara *et al.* 2006a; Table 1) and after PM (TeKrony *et al.* 1979, 1980; Gibson & Mullen 1996; Hampton 2000). Shinohara *et al.* (2008) examined the relationship between vigour test results for 262 garden pea seeds lots produced in New Zealand and climate data in five regions over four consecutive production seasons, and while regional and seasonal variation for vigour occurred, these variations were significantly associated with temperature during seed development—generally the higher the temperature, the lower the seed vigour.

The susceptibility of seeds to loss of vigour following high-temperature stress depends on the stage of development (Shinohara *et al.* 2006a). Using hollow heart, a physiological disorder of germinating pea seeds (Halligan 1986) as a seed vigour indicator (Castillo *et al.* 1993), Shinohara *et al.* (2006a) found in a field study that hourly thermal time (HTT, measured in degree hours, °Ch, T<sub>b</sub>=25 °C) when seeds were at the green-wrinkled pod stage (700–800 mg/g seed

Table 3. Effect of temperature during seed development and maturation on nucleotide content, mitochondrial respiration rate and adenylate energy charge (AEC) of excised wheat embryos after 4 h imbibition (adapted from Grass & Burris 1995)

|   | Temperature regime* |          |          |
|---|---------------------|----------|----------|
|   | 20/15 °C            | 28/21 °C | 36/29 °C |
| Nucleotide content†<br>(nmol/g)   |                     |          |          |
| AMP   | 400                 | 370      | 448      |
| ADP   | 571                 | 518      | 412      |
| ATP   | 452                 | 393      | 141      |
| Total   | 1422                | 1180     | 1001     |
| State 3 mitochondrial<br>respiration rate<br>(nmol O <sub>2</sub> /min) | 9.7                 | 6.9      | 5.3      |
| AEC‡  | 0.52                | 0.51     | 0.35     |

\* Day/night with 8 h day temperature and 16 h night temperature.

† AMP, adenosine monophosphate; ADP, adenosine diphosphate; ATP, adenosine triphosphate.

‡ AEC expressed as the ratio (ATP+0.5ADP/ATP+ADP+AMP)=energy status of the seed (Atkinson 1968).

moisture content (SMC)) was significantly correlated with hollow heart incidence at harvest, and that 100 °Ch were required to induce the condition. There was no such relationship between HTT and hollow heart after PM. While there were cultivar differences, for one cultivar there was a linear increase in hollow heart incidence as the degree hours (°Ch) increased. In a follow-up controlled environment study, Shinohara *et al.* (2006b) confirmed this result, by demonstrating that exposure to day/night temperatures of 30 and 25 °C, respectively, for 4 days (240 °Ch, Tb=25 °C) at the green-wrinkled pod stage induced hollow heart, but exposure to the same conditions at the beginning of seed fill (>800 mg/g SMC), PM (550–650 mg/g SMC) or after PM did not (Table 2). Single-seed conductivity (which is an indicator of cell membrane integrity – see Powell 2006) was increased only after exposure of the developing seeds to the high temperature at or after PM, and not before (Table 1).

Seed vigour loss is associated with seed physiological deterioration (Powell 1988; Hampton & Coolbear 1990), and lipid peroxidation is the most frequently cited cause (McDonald 1999). Lipid peroxidation causes cellular degeneration through free radical assault on important cellular molecules and structures (Wilson & McDonald 1986). McDonald (1999), in his model of seed deterioration, proposed four types of cell damage, viz. mitochondrial dysfunction, enzyme

Table 4. Effect of sowing date at the same field site on HTT (Tb=25 °C) and the number of hours of exposure to temperature exceeding 25 and 30 °C during the period when SMC was between 700 and 800 mg/g for garden pea cvar Alderman (adapted from Shinohara *et al.* 2006a).

| Sowing date | SMC 700–800 mg/g |           |                     |        | PM*    |
|-------------|------------------|-----------|---------------------|--------|--------|
|             | Period†          | HTT (°Ch) | No. of hours >25 °C | >30 °C |        |
| 26 Sep      | 31 Dec–4 Jan     | 198       | 45                  | 19     | 11 Jan |
| 21 Oct      | 8 Jan–13 Jan     | 106       | 38                  | 8      | 21 Jan |
| 19 Nov      | 1 Feb–7 Feb      | 21        | 5                   | 0      | 18 Feb |

\* Dates when seeds were adjudged to have reached PM.

† Period when SMC was between 700 and 800 mg/g.

inactivation, membrane degradation and genetic damage.

Grass & Burris (1995) reported that high-temperature stress of the parent plant caused mitochondrial degeneration and reduced adenosine triphosphate (ATP) accumulation, energy levels and rates of oxygen uptake in imbibing wheat embryos (Table 3), providing clear evidence for metabolic changes at the mitochondrial level in early seed germination in response to heat stress during seed development and maturation. High temperatures during reproductive growth increase seed cell membrane damage (Nilsen & Orcutt 1996; Shinohara *et al.* 2006b) so that electrolyte leakage from seeds is increased (Castillo *et al.* 1994; Spears *et al.* 1997; Shinohara *et al.* 2006b). High leachate conductivity in pea has been associated with dead/deteriorating tissue on the abaxial surface of the cotyledons (Powell 1985; Shinohara *et al.* 2006b), and on the adaxial cotyledonary surface for hollow heart (Don *et al.* 1984; Shinohara *et al.* 2006b). However, temperature stress also results in damage to the shoot apical meristem of the embryonic axis (Fu *et al.* 1988; Senaratna *et al.* 1988). Membrane disorganization would reduce mitochondrial efficiency and may allow the release of peroxidative enzymes capable of causing subsequent cellular damage after imbibition has begun (McDonald 1999).

If heat stress leads to mitochondrial dysfunction and membrane damage, it may also result in reduced enzyme activity (e.g. decreased  $\alpha$ -amylase – Shephard *et al.* 1996) and genetic damage (e.g. decreased DNA synthesis – Cruz-Garcia *et al.* 1995). Whether these



and other seed deteriorative changes (see McDonald 1999) occur following heat stress during seed development and maturation is yet to be determined.

## CONCLUSIONS

The environment during seed development and maturation can significantly reduce seed quality (Dornbos 1995; Gusta *et al.* 2004; Egli *et al.* 2005; Shinohara *et al.* 2008), particularly seed vigour. How likely is it that elevated CO<sub>2</sub> levels and temperature increases of up to 3 °C by 2050 will further increase this loss of seed quality? To answer this question accurately will require substantially more research in order to determine the critical periods during seed development when seeds are sensitive to environmental stresses, and for temperature, how this interacts with the duration of exposure to elevated temperatures which are deleterious to seed quality. For example, Shinohara *et al.* (2006b) found that during the rapid seed filling stage in pea, a temperature of 30/25 °C for 2 days (120 °Ch at Tb=25 °C) did not induce hollow heart, but 4 days induced hollow heart in one cultivar (see Table 2), and 6 days (360 °Ch) induced the condition in both cultivars used (0.43 in cvar Alderman and 0.23 in cvar Early Onward).

From the information that is available, it can be concluded that predicted environmental changes will lead to the increased occurrence of loss of seed quality, particularly seed vigour and possibly germination. While seed mass will also change, this does not necessarily imply any negative effect on germination or vigour. To minimize the risk of reductions in seed quality the seed industry will therefore have to consider:

- (a) Moving seed production to the limits of adaptation either in latitude (northern or southern) or in elevation (highland and mountainous regions) in order to reduce the chances of environmental stress (Egli *et al.* 2005; Shinohara *et al.* 2008).
- (b) Changing sowing date so that seed filling occurs at lower temperatures (Castillo *et al.* 1994; Egli *et al.* 2005; Shinohara *et al.* 2006a). The latter authors demonstrated that for the pea cultivar Alderman, HTT (Tb=25 °C) during the rapid seed filling stage was 198, 106 and 21 °Ch for sowings at the same site in September, October and November, respectively, and the number of hours during this stage when temperature exceeded 25 or 30 °C also reduced as sowing date was delayed (Table 4).

- (c) Exploring genotypic differences in the ability to acquire and retain good seed quality in stressful environments, firstly among existing cultivars (Spears *et al.* 1997; Shinohara *et al.* 2006a), and in the breeding of new cultivars (Ainsworth *et al.* 2008b).

## REFERENCES

- AINSWORTH, E. A., ROGERS, A. & LEAKEY, A. D. B. (2008a). Targets for crop biotechnology in a future high-CO<sub>2</sub> and high-O<sub>3</sub> world. *Plant Physiology* **147**, 13–19.
- AINSWORTH, E. A., BEIER, C., CALFAPIETRA, C., CEULEMANS, R., DURAND-TARDIF, M., FARQUHAR, G. D., GODBOLD, D. L., HENDREY, G. R., HICKLER, T., KADUK, J., KARNOSKY, D. F., KIMBALL, B. A., KÖRNER, C., KOORNNEEF, M., LAFARGE, T., LEAKEY, A. D., LEWIN, K. F., LONG, S. P., MANDERSCHIED, R., MCNEIL, D. L., MIES, T. A., MIGLIETTA, F., MORGAN, J. A., NAGY, J., NORBY, R. J., NORTON, R. M., PERCY, K. E., ROGERS, A., SOUSSANA, J. F., STITT, M., WEIGEL, H. J. & WHITE, J. W. (2008b). Next generation of elevated CO<sub>2</sub> experiments with crops: a critical investment for feeding the future world. *Plant, Cell and Environment* **31**, 1317–1324.
- ALMEKINDERS, C. J. M. & LOUWAARS, N. P. (1999). *Farmer's Seed Production. New Approaches and Practices*. London: ITDG Publishing.
- ANDALO, C., GODELLE, B., LEFRANC, M., MOUSSEAU, M. & TILL-BOTTRAUD, I. (1996). Elevated CO<sub>2</sub> decreases seed germination in *Arabidopsis thaliana*. *Global Change Biology* **2**, 129–135.
- ALLEN, L. H. Jr & BOOTE, K. J. (2000). Crop ecosystem responses to climatic change: soybean. In *Climate Change and Global Crop Productivity* (Eds K. R. Reddy & H. F. Hodges), pp. 133–160. Wallingford, UK: CAB International.
- ATKINSON, D. E. (1968). The energy charge of the adenylate pool as a regulatory parameter. *Biochemistry* **7**, 4030–4034.
- BAKER, J. T., ALLEN, L. H. Jr, BOOTE, K. J., JONES, P. & JONES, J. W. (1989). Response of soybean to air temperature and carbon dioxide concentration. *Crop Science* **29**, 98–105.
- BRUINS, M. (2009). The evolution and contribution of plant breeding to global agriculture. In *Proceedings of the Second World Seed Conference: Responding to the Challenges of a Changing World: The Role of New Plant Varieties and High Quality Seed in Agriculture*, pp. 18–31. Rome: FAO.
- CASTILLO, A. G., HAMPTON, J. G. & COOLBEAR, P. (1993). Effect of population density on within canopy environment and seed vigour in garden pea (*Pisum sativum* L.). *Proceedings of the Annual Conference of the Agronomy Society of New Zealand* **23**, 99–106.
- CASTILLO, A. G., HAMPTON, J. G. & COOLBEAR, P. (1994). Effect of sowing date and harvest timing on seed vigour in garden pea (*Pisum sativum* L.). *New Zealand Journal of Crop and Horticultural Science* **22**, 91–95.

- CASTRO, J., HODAR, J. A. & GOMEZ, J. M. (2006). Seed size. In *Handbook of Seed Science and Technology* (Ed. A. S. Basra), pp. 397–427. New York: Food Products Press.
- CECCARELLI, S., GRANDO, S., MAATOUGUI, M., MICHAEL, M., SLASH, M., HAGHPARAST, R., RAHMANIAN, M., TAHERI, A., AL-YASSIN, A., BENBELKACEM, A., LABDI, M., MIMOUN, H. & NACHIT, M. (2010). Plant breeding and climate changes. *Journal of Agricultural Science, Cambridge* **148**, 627–637.
- COOLBEAR, P. (1995). Mechanisms of seed deterioration. In *Seed Quality: Basic Mechanisms and Agricultural Implications* (Ed. A. S. Basra), pp. 223–277. New York: Food Products Press.
- CRUZ-GARCIA, F., GONZALEZ-HERNANDEZ, V. A., MOLINA-MORENO, J. & VAZQUEZ-RAMOS, J. M. (1995). Seed deterioration and respiration as related to DNA metabolism in germinating maize. *Seed Science and Technology* **23**, 477–486.
- DIJKSTRA, P., SCHAPENDONK, H. M. C., GROENWOLD, K., JANSEN, M. & VAN DE GEIJN, S. C. (1999). Seasonal changes in the response of winter wheat to elevated atmospheric CO<sub>2</sub> concentration grown in open-top chambers and field tracking enclosures. *Global Change Biology* **5**, 563–576.
- DON, R., BUSTAMANTE, L., RENNIE, W. J. & SEDDON, M. G. (1984). Hollow heart in pea (*Pisum sativum* L.). *Seed Science and Technology* **12**, 707–721.
- DORNBOS, D. L. Jr (1995). Production environment and seed quality. In *Seed Quality: Basic Mechanisms and Agricultural Implications* (Ed. A. S. Basra), pp. 119–152. New York: Food Products Press.
- DORNBOS, D. L. Jr & McDONALD, M. B. (1986). Mass and composition of developing soybean seeds at five reproductive growth stages. *Crop Science* **26**, 624–630.
- EDWARDS, G. R., CLARK, H. & NEWTON, P. C. D. (2001). The effects of elevated CO<sub>2</sub> on seed production and seedling recruitment in a sheep-grazed pasture. *Oecologia* **127**, 383–394.
- EGLI, D. B., TE KRONY, D. M., HEITHOLT, J. J. & RUPE, J. (2005). Air temperature during seed filling and soybean seed germination and vigour. *Crop Science* **45**, 1329–1335.
- ESASHI, Y., OOSHIMA, Y., MICHIHARA AABE, M., KUROTA, A. & SATOH, S. (1986). CO<sub>2</sub>-enhanced C<sub>2</sub>H<sub>4</sub> production in tissues of imbibed cocklebur seeds. *Australian Journal of Plant Physiology* **13**, 417–429.
- FARNSWORTH, E. J. & BAZAZ, F. A. (1995). Inter- and intra-generic differences in growth, reproduction, and fitness of nine herbaceous annual species grown in elevated CO<sub>2</sub> environments. *Oecologia* **104**, 454–466.
- FENNER, M. (1991). The effects of the parent environment on seed germinability. *Seed Science Research* **1**, 75–84.
- FU, J. R., LU, X. H., CHEN, R. Z., ZHANG, B. Z., LIU, Z. S., LI, Z. S. & CAI, C. Y. (1988). Osmoconditioning of peanut (*Arachis hypogaea* L.) seeds with PEG to improve vigour and some biochemical activities. *Seed Science and Technology* **16**, 197–212.
- GIBSON, L. R. & MULLEN, R. E. (1996). Soybean seed quality reductions by high day and night temperature. *Crop Science* **36**, 1615–1619.
- GIBSON, L. R. & PAULSEN, G. M. (1999). Yield components of wheat grown under high temperature stress during reproductive growth. *Crop Science* **39**, 1841–1846.
- GORNALL, J., BETTS, R., BURKE, E., CLARK, R., CAMP, J., WILLETT, K. & WILTSHIRE, A. (2010). Implications of climate change for agricultural productivity in the early twenty-first century. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 2973–2989.
- GRASS, L. & BURRIS, J. S. (1995). Effect of heat stress during seed development and maturation on wheat (*Triticum durum*) seed quality. II. Mitochondrial respiration and nucleotide pools during early germination. *Canadian Journal of Plant Science* **75**, 831–839.
- GREEN, D. E., PINNELL, E. L., CAVANAUGH, L. E. & WILLIAMS, L. F. (1965). Effect of planting date and maturity date on soybean seed quality. *Agronomy Journal* **57**, 165–168.
- GUSTA, L. V., JOHNSON, E. N., NESBITT, N. T. & KIRKLAND, K. J. (2004). Effect of seeding date on canola seed quality and seed vigour. *Canadian Journal of Plant Science* **84**, 463–471.
- GUTTERMAN, Y. (2000). Maternal effects on seed during development. In *Seeds: The Ecology of Regeneration in Plant Communities* (Ed. M. Fenner), pp. 59–84. Wallingford, UK: CAB International.
- HALLIGAN, E. A. (1986). The effect of elevated temperatures and their duration on the incidence of hollow heart in pea seeds. *Annals of Applied Biology* **109**, 619–625.
- HAMPTON, J. G. (2000). Producing quality seed: the problem of seed vigour. In *Current Research on Seeds in New Zealand* (Eds M. T. McManus, H. A. Outred & K. M. Pollock), pp. 53–67. Agronomy Society of New Zealand Special Publication No. 12. Christchurch, New Zealand: Agronomy Society of New Zealand.
- HAMPTON, J. G. (2002). What is seed quality? *Seed Science and Technology* **30**, 1–10.
- HAMPTON, J. G. & COOLBEAR, P. (1990). Potential versus actual seed performance – can vigour testing provide an answer? *Seed Science and Technology* **18**, 215–228.
- HARA, Y. & TORIYAMA, K. (1998). Seed nitrogen accelerates the rates of germination, emergence and establishment of rice plants. *Soil Science and Plant Nutrition* **44**, 359–366.
- HE, J. S., FLYNN, D. F. B., WOLFE-BELLIN, K., FANG, J. & BAZAZ, F. A. (2005). CO<sub>2</sub> and nitrogen, but not population density, alter the size and C/N ratio of *Phytolacca americana* seeds. *Functional Ecology* **19**, 437–444.
- HIKOSAKA, K., KINUGASA, T., OIKAWA, S., ONODA, Y. & HIROSE, T. (2011). Effects of elevated CO<sub>2</sub> concentration on seed production in C<sub>3</sub> annual plants. *Journal of Experimental Botany* **62**, 1523–1530.
- HOUGHTON, J. T., DING, Y., GRIGGS, D. J., NOGUER, M., VAN DER LINDEN, P. J., DAI, X., MASKELL, K. & JOHNSON, C. A. (2001). *Climate Change 2001: The Scientific Basis. Report of the IPCC*. Cambridge, UK: Cambridge University Press.
- HUXMAN, T. E., HAMERLYNCK, E. P., JORDAN, D. N., SALSMAN, K. A. & SMITH, S. D. (1998). The effects of parental CO<sub>2</sub> environment on seed quality and subsequent seedling performance in *Bromus rubens*. *Oecologia* **114**, 207–208.
- ISTA (2011). *International Rules for Seed Testing*. Bassersdorf, Switzerland: International Seed Testing Association.
- JABLONSKI, L. M., WANG, X. & CURTIS, P. S. (2002). Plant reproduction under elevated CO<sub>2</sub> conditions: a meta-analysis of reports. *New Phytologist* **156**, 9–26.

- JAGGARD, K. W., QI, A. & OBER, E. S. (2010). Possible changes to arable crop yields by 2050. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 2835–2851.
- KHALIL, S. K., MEXAL, J. G. & MURRAY, L. W. (2001). Soybean seed matured on different dates affect seed quality. *Pakistan Journal of Biological Sciences* **4**, 365–370.
- KHALIL, S. K., MEXAL, J. G., REHMAN, A., KHAN, A. Z., WAHAB, S., ZUBAIR, M., KHALIL, I. H. & MOHAMMAD, F. (2010). Soybean mother plant exposure to temperature stress and its effect on germination under osmotic stress. *Pakistan Journal of Botany* **42**, 213–225.
- LEUBNER-METZGER, G. (2006). Hormonal interactions during seed dormancy release and germination. In *Handbook of Seed Science and Technology* (Ed. A. S. Basra), pp. 303–342. New York: Food Products Press.
- MCDONALD, M. B. (1999). Seed deterioration: physiology, repair and assessment. *Seed Science and Technology* **27**, 177–237.
- MCDONALD, M. B. & NELSON, C. J. (1986). *Physiology of Seed Deterioration*. Special Publication No. 11, Madison, WI, USA: Crop Science Society of America.
- MIYAGI, K. M., KINUGASA, T., HIKOSAKA, K. & HIROSE, T. (2007). Elevated CO<sub>2</sub> concentration, nitrogen use, and seed production in annual plants. *Global Change Biology* **13**, 2161–2170.
- MUSGRAVE, M. E., STRAIN, B. R. & SIEDOW, J. N. (1986). Response of two pea hybrids to CO<sub>2</sub> enrichment: a test of the energy overflow hypothesis for alternative respiration. *Proceedings of the National Academy of Sciences, USA* **83**, 8157–8161.
- NILSEN, E. T. & ORCUTT, D. M. (1996). Plant membranes as environmental sensors. In *The Physiology of Plants Under Stress: Abiotic Factors* (Eds E. T. Nilsen & D. M. Orcutt), pp. 50–82. New York: John Wiley and Sons.
- PELTONEN-SAINIO, P., JAUHIAINEN, L. & HAKALA, K. (2011). Crop responses to temperature and precipitation according to long-term multi-location trials at high-latitude conditions. *Journal of Agricultural Science, Cambridge* **149**, 49–62.
- POWELL, A. A. (1985). Impaired membrane integrity: a fundamental cause of seed quality differences in peas. In *The Pea Crop: a Basis for Improvement* (Eds P. D. Hebblethwaite, M. C. Heath & T. C. K. Dawkins), pp. 383–394. London: Butterworths.
- POWELL, A. A. (1988). Seed vigour and field establishment. *Advances in Research and Technology of Seeds* **11**, 29–61.
- POWELL, A. A. (2006). Seed vigour and its assessment. In *Handbook of Seed Science and Technology* (Ed. A. S. Basra), pp. 603–648. New York: Food Products Press.
- PRASAD, P. V. V., BOOTE, K. J., ALLEN, L. H. JR & THOMAS, J. M. G. (2002). Effects of elevated temperature and carbon dioxide on seed-set and yield of kidney bean (*Phaseolus vulgaris* L.). *Global Change Biology* **8**, 710–721.
- QUADERI, M. M. & REID, D. M. (2005). Growth and physiological responses of canola (*Brassica napus*) to UV-B and CO<sub>2</sub> under controlled environment conditions. *Physiologia Plantarum* **125**, 247–259.
- SENARATNA, T., GUSSE, J. F. & MCKERSIE, B. D. (1988). Age-induced changes in cellular membranes of imbibed soybean axes. *Physiologia Plantarum* **73**, 85–91.
- SHEPHARD, H. L., NAYLOR, R. E. L. & STUCHBURY, T. (1996). The influence of seed maturity at harvest and drying method on the embryo,  $\alpha$ -amylase activity and seed vigour in sorghum (*Sorghum bicolor* (L.) Moench). *Seed Science and Technology* **24**, 245–249.
- SHINOHARA, T., HAMPTON, J. G. & HILL, M. J. (2006a). Effects of the field environment before and after seed physiological maturity on hollow heart occurrence in garden pea (*Pisum sativum*). *New Zealand Journal of Crop and Horticultural Science* **34**, 247–255.
- SHINOHARA, T., HAMPTON, J. G. & HILL, M. J. (2006b). Location of deterioration within garden pea (*Pisum sativum*) cotyledons is associated with the timing of exposure to high temperature. *New Zealand Journal of Crop and Horticultural Science* **34**, 299–309.
- SHINOHARA, T., HAMPTON, J. G., HILL, M. J., JUNTAKOOL, S., SUPRAKARN, S. & SAGWANSUPYAKORN, C. (2008). Variations in pea (*Pisum sativum* L.) seed vigour among regions of production and cropping seasons are associated with temperature during reproductive growth. *Journal of the Japanese Society of Agricultural Technology Management* **14**, 148–155.
- SMITH, S. D., HUXMAN, T. E., ZITZER, S. F., CHARLET, T. N., HOUSMAN, D. C., COLEMAN, J. S., FENSTERMAKER, L. K., SEEMANN, J. R. & NOWAK, R. W. (2000). Elevated CO<sub>2</sub> increases productivity and invasive species success in an arid ecosystem. *Nature* **408**, 79–82.
- SOLOMON, S., QIN, D., MANNING, M., CHEN, Z., MARQUIS, M., AVERYT, K. B., TIGNOR, M. & MILLER, H. L. (2007). Technical summary. In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Annual Report of the Intergovernmental Panel on Climate Change* (Eds S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor & H. L. Miller), pp. 19–840. Cambridge, UK: Cambridge University Press.
- SPEARS, J. F., TEKRONY, D. M. & EGLI, D. B. (1997). Temperature during seed filling and soybean seed germination and vigour. *Seed Science and Technology* **25**, 233–244.
- STEINGER, T., GALL, R. & SCHMID, B. (2000). Maternal and direct effects of elevated CO<sub>2</sub> on seed provisioning, germination and seedling growth in *Bromus erectus*. *Oecologia* **123**, 475–480.
- TEKRONY, D. M., EGLI, D. B., BALLES, J., PFEIFFER, T. & FELLOWS, R. J. (1979). Physiological maturity in soybeans. *Agronomy Journal* **71**, 771–775.
- TEKRONY, D. M., EGLI, D. B. & BALLES, J. (1980). The effects of the field production environment on soybean seed quality. In *Seed Production* (Ed. P. D. Hebblethwaite), pp. 403–426. London: Butterworth and Co Ltd.
- THOMAS, J. M. G., PRASAD, P. V. V., BOOTE, K. J. & ALLEN, L. H. (2009). Seed composition, seedling emergence and early seedling vigour of red kidney bean seed produced at elevated temperature and carbon dioxide. *Journal of Agronomy and Crop Science* **195**, 148–156.
- WAGNER, J., LUSCHER, A., HILLEBRAND, C., KOBALD, B., SPITALER, N. & LARCHER, W. (2001). Sexual reproduction of *Lolium perenne* L. and *Trifolium repens* L. under free air CO<sub>2</sub>



- enrichment (FACE) at two levels of nitrogen application. *Plant, Cell and Environment* **24**, 957–965.
- WAY, D. A., LADEAU, S. L., MCCARTHY, H. R., CLARK, J. S., OREN, R., FINZI, A. C. & JACKSON, R. B. (2010). Greater seed production in elevated CO<sub>2</sub> is not accompanied by reduced seed quality in *Pinus taeda* L. *Global Change Biology* **16**, 1046–1056.
- WEIGAND, C. L. & CUELLER, J. A. (1981). Duration of grain filling and kernel weight of wheat as affected by temperature. *Crop Science* **21**, 95–101.
- WHEELER, T. R., CRAUFURD, P. Q., ELLIS, R. H., PORTER, J. R. & PRASAD, P. V. V. (2000). Temperature variability and the yield of annual crops. *Agriculture, Ecosystems and Environment* **82**, 159–167.
- WILSON, D. O. & McDONALD, M. B. (1986). The lipid peroxidation model of seed deterioration. *Seed Science and Technology* **14**, 269–300.
- WULF, R. D. & ALEXANDER, H. M. (1985). Intraspecific variation in the response to CO<sub>2</sub> enrichment in seeds and seedlings of *Plantago lanceolata*. *Oecologia* **66**, 458–460.
- YOUNG, L. W., WILEN, R. W. & BONHAM-SMITH, P. C. (2004). High temperature stress of *Brassica napus* during flowering reduces micro-and megagametophyte fertility, induces fruit abortion, and disrupts seed production. *Journal of Experimental Botany* **55**, 485–495.
- ZISKA, L. H. & BUNCE, J. A. (1993). The influence of elevated CO<sub>2</sub> and temperature on seed germination and emergence from soil. *Field Crops Research* **34**, 147–157.